ELSEVIER

Molecular Brain Research 75 (2000) 355–357

MOLECULAR BRAIN RESEARCH

Author Index

Ahn, K.Y., see Lee, Z.H. (75) 237
Ambrosini, A., Tininini, S., Barassi, A., Racagni, G., Sturani, E. and Zippel, R. cAMP cascade leads to Ras activation in cortical neurons (75) 54
Ambrosio, S., see Goren, N. (75) 138
Anisman, H., see Plata-Salamán, C.R. (75) 248
Arancibia, S., see Givalois, L. (75) 166
Arimura, A., see Zhou, C.J. (75) 150

Bae, C.S., see Lee, Z.H. (75) 237
Bae, H.R., see Lee, S.A. (75) 16
Bae, K.W., see Lee, S.A. (75) 16
Barassi, A., see Ambrosini, A. (75) 54
Bartrons, R., see Goren, N. (75) 138
Bayliss, D.A., see Talley, E.M. (75) 159
Bechtold, D.A. and Brown, I.R. Heat shock proteins Hsp27 and Hsp32 localize to synaptic sites in the rat cerebellum following hyperthermia (75) 309
Bernabeu, R., Di Scala, G. and Zwiller, J. Odor regulates the expression of the mitogen-activated protein kinase phosphatase gene *hVH-5* in bilateral entorhinal cortex-lesioned rats (75) 113
Blank, T., see Radulovic, J. (75) 271
Bond, B.C., see Harrison, D.C. (75) 143
Bowery, N.G., see Ingram, E.M. (75) 96
Bramham, C., see Valentine, G. (75) 337
Broberger, C., Nylander, I., Geijer, T., Terenius, L., Hökfelt, T. and Georgieva, J. Differential effects of intrastriatally infused fully and endcap phosphorothioate antisense oligonucleotides on morphology, histochemistry and prodynorphin expression in rat brain (75) 25
Brown, I.R., see Bechtold, D.A. (75) 309
Bureau, Y., see Plata-Salamán, C.R. (75) 248
Burke, R.E., see Kholodilov, N.G. (75) 281

Campbell, C.A., see Harrison, D.C. (75) 143
Cavallaro, S., see D'Agata, V. (75) 345
Chakravarty, S., see Valentine, G. (75) 337
Chan, S.L., see Gary, D.S. (75) 89
Chen, N. and Justice Jr., J.B. Differential effect of structural modification of human dopamine transporter on the inward and outward transport of dopamine (75) 208
Christakos, S., see Gary, D.S. (75) 89
Coker, T.N., see Scammell, J.G. (75) 8
Corbitt, J., see Sands, S.A. (75) 1
Craft, C.M. and Zhan-Poe, X. Identification of specific histidine residues and the carboxyl terminus are essential for serotonin *N*-acetyltransferase enzymatic activity (75) 198
Cuvelier, L., see Depaepe, V. (75) 76

D'Agata, V., Zhao, W. and Cavallaro, S. Cloning and distribution of the rat parkin mRNA (75) 345
Damm, A., see Mielke, K. (75) 128
David Clark, J., see Li, X. (75) 179
Davis, R.P., see Harrison, D.C. (75) 143
De Kloet, E.R., see Schaaf, M.J.M. (75) 342
Depaepe, V., Cuvelier, L., Thöny, B. and Résibois, A. Pterin-4α-carbinolamine dehydratase in rat brain. I. Patterns of co-localization with tyrosine hydroxylase (75) 76
Depaulis, A., see Talley, E.M. (75) 159
Di Scala, G., see Bernabeu, R. (75) 113
Dixon, W.T., see Weaver, S.A. (75) 46
Doi, T., Ueda, Y., Tokumaru, J., Mitsuyama, Y. and Willmore, L.J. Sequential changes in glutamate transporter mRNA levels during Fe^{3+}-induced epileptogenesis (75) 105
Durston, A.J., see Holling, T.M. (75) 70
Duurland, R., see Schaaf, M.J.M. (75) 342

Eggold, J.R., see Naeve, G.S. (75) 185
Emson, P.C., see Ingram, E.M. (75) 96
Ernfors, P., see Linnarsson, S. (75) 61

Fernald, R.D., see Latimer, V.S. (75) 287
Flynn, M.C., see Plata-Salamán, C.R. (75) 248
Forsayeth, J., see Sweileh, W. (75) 293
Foster, A.C., see Naeve, G.S. (75) 185

Gale, K., see Kondratyev, A. (75) 216
Gary, D.S., Sooy, K., Chan, S.L., Christakos, S. and Mattson, M.P. Concentration- and cell type-specific effects of calbindin D28k on vulnerability of hippocampal neurons to seizure-induced injury (75) 89
Garyfallou, V.T., see Latimer, V.S. (75) 287
Gayle, D., see Plata-Salamán, C.R. (75) 248
Geijer, T., see Broberger, C. (75) 25
Georgieva, J., see Broberger, C. (75) 25
Givalois, L., Arancibia, S. and Tapia-Arancibia, L. Concomitant changes in CRH mRNA levels in rat hippocampus and hypothalamus following immobilization stress (75) 166
Goren, N., Manzano, A., Riera, L., Ambrosio, S., Ventura, F. and Bartrons, R. 6-Phosphofructo-2-kinase/fructose-2,6-bisphosphatase expression in rat brain during development (75) 138
Green, C.B., see Zhu, H. (75) 303

Hardy, S., see Sweileh, W. (75) 293
Harrison, D.C., Medhurst, A.D., Bond, B.C., Campbell, C.A., Davis, R.P. and Philpott, K.L. The use of quantitative RT-PCR to measure mRNA expression in a rat model of focal ischemia — caspase-3 as a case study (75) 143
Herdegen, T., see Mielke, K. (75) 128
Herkenham, M., see Valentine, G. (75) 337
Hirabayashi, T., see Zhou, C.J. (75) 150
Hökfelt, T., see Broberger, C. (75) 25
Holling, T.M., Van Herp, F., Durston, A.J. and Martens, G.J.M. Differential onset of expression of mRNAs encoding proopiomelanocortin, prohormone convertases 1 and 2, and granin family members during *Xenopus laevis* development (75) 70
Huey, K.A. and Powell, F.L. Time-dependent changes in dopamine D_2-receptor mRNA in the arterial chemoreflex pathway with chronic hypoxia (75) 264
Hung, H.-C., Tsai, M.J., Wu, H.C. and Lee, E.H.Y. Age-dependent increase in *C7-1* gene expression in rat frontal cortex (75) 330
Hutson, S.M., see Kholodilov, N.G. (75) 281

Ilyin, S.E., see Plata-Salamán, C.R. (75) 248
Ingram, E.M., Tessler, S., Bowery, N.G. and Emson, P.C. Glial glutamate transporter mRNAs in the genetically absence epilepsy rat from Strasbourg (75) 96
Inoue, M., see Renbäck, K. (75) 350

Justice Jr., J.B., see Chen, N. (75) 208

Kammermeier, J., see Radulovic, J. (75) 271
Kang, E.K., see Lee, S.A. (75) 16
Kelly, M.E., see Plata-Salamán, C.R. (75) 248
Kholodilov, N.G., Neystat, M., Oo, T.F., Hutson, S.M. and Burke, R.E.
Upregulation of cytosolic branched chain aminotransferase in substantia nigra following developmental striatal target injury (75) 281
Kikuyama, S., see Zhou, C.J. (75) 150
Kim, H.-H., see Lee, Z.H. (75) 237
Kim, J.K., see Lee, Z.H. (75) 237
Kim, K.K., see Lee, Z.H. (75) 237
Kohama, S.G., see Latimer, V.S. (75) 287
Kojima, T., Nakajima, K. and Mikoshiba, K.
The *disabled 1* gene is disrupted by a replacement with L1 fragment in *yotari* mice (75) 121
Kondo, H., see Kudo, M. (75) 172
Kondratyev, A. and Gale, K.
Intracerebral injection of caspase-3 inhibitor prevents neuronal apoptosis after kainic acid-evoked status epilepticus (75) 216
Kudo, M., Saito, S., Owada, Y., Suzaki, H. and Kondo, H.
Localization of mRNA for SHIP2, SH2 domain-containing inositol polyphosphate 5-phosphatase, in the brain of developing and mature rats (75) 172

LaRue, S., see Zhu, H. (75) 303
Latimer, V.S., Rodrigues, S.M., Garyfallou, V.T., Kohama, S.G., White, R.B., Fernald, R.D. and Urbanski, H.F.
Two molecular forms of gonadotropin-releasing hormone (GnRH-I and GnRH-II) are expressed by two separate populations of cells in the rhesus macaque hypothalamus (75) 287
Lee, E.H.Y., see Hung, H.-C. (75) 330
Lee, S.A., Park, J.K., Kang, E.K., Bae, H.R., Bae, K.W. and Park, H.T.
Calmodulin-dependent activation of p38 and p42/44 mitogen-activated protein kinases contributes to c-*fos* expression by calcium in PC12 cells: modulation by nitric oxide (75) 16
Lee, Z.H., Kim, H.-H., Ahn, K.Y., Seo, K.H., Kim, J.K., Bae, C.S. and Kim, K.K.
Identification of a brain specific protein that associates with a Refsum disease gene product, phytanoyl-CoA alpha-hydroxylase (75) 237
Lemaigre-Dubreuil, Y., see Vernet-der Garabedian, B. (75) 259
Li, X. and David Clark, J.
Chronic morphine exposure and the expression of heme oxygenase type 2 (75) 179
Ling, N., see Naeve, G.S. (75) 185
Linnarsson, S., Willson, C.A. and Ernfors, P.
Cell death in regenerating populations of neurons in BDNF mutant mice (75) 61
Loring, R.H., see Sweileh, W. (75) 293

Manzano, A., see Goren, N. (75) 138
Mariani, J., see Vernet-der Garabedian, B. (75) 259
Martens, G.J.M., see Holling, T.M. (75) 70
Mattson, M.P., see Gary, D.S. (75) 89
McIntyre, D.C., see Plata-Salamán, C.R. (75) 248
Medhurst, A.D., see Harrison, D.C. (75) 143
Middendorff, R., see Müller, D. (75) 321
Mielke, K., Damm, A., Yang, D.D. and Herdegen, T.
Selective expression of JNK isoforms and stress-specific JNK activity in different neural cell lines (75) 128
Mikoshiba, K., see Kojima, T. (75) 121
Mitsuyama, Y., see Doi, T. (75) 105
Morilak, D.A., see Sands, S.A. (75) 1
Mukhopadhyay, A.K., see Müller, D. (75) 321
Müller, D., Olcese, J., Mukhopadhyay, A.K. and Middendorff, R.
Guanylyl cyclase-B represents the predominant natriuretic peptide receptor expressed at exceptionally high levels in the pineal gland (75) 321

Naeve, G.S., Vana, A.M., Eggold, J.R., Verge, G., Ling, N. and Foster, A.C.
Expression of rat insulin-like growth factor binding protein-6 in the brain, spinal cord, and sensory ganglia (75) 185
Nakajima, K., see Kojima, T. (75) 121
Nakajo, S., see Zhou, C.J. (75) 150
Neystat, M., see Kholodilov, N.G. (75) 281
Nijholt, I., see Radulovic, J. (75) 271
Nikolopoulos, S.N., see Scammell, J.G. (75) 8
Nyberg, F., see Renbäck, K. (75) 350
Nylander, I., see Broberger, C. (75) 25

Olcese, J., see Müller, D. (75) 321
Oo, T.F., see Kholodilov, N.G. (75) 281
Owada, Y., see Kudo, M. (75) 172

Park, H.T., see Lee, S.A. (75) 16
Park, J.K., see Lee, S.A. (75) 16
Perez-Reyes, E., see Talley, E.M. (75) 159
Philpott, K.L., see Harrison, D.C. (75) 143
Plata-Salamán, C.R., Ilyin, S.E., Turrin, N.P., Gayle, D., Flynn, M.C., Romanovitch, A.E., Kelly, M.E., Bureau, Y., Anisman, H. and McIntyre, D.C.
Kindling modulates the IL-1β system, TNF-α, TGF-β1, and neuropeptide mRNAs in specific brain regions (75) 248
Powell, F.L., see Huey, K.A. (75) 264

Racagni, G., see Ambrosini, A. (75) 54
Radulovic, J., Blank, T., Nijholt, I., Kammermeier, J. and Spiess, J.
In vivo NMDA/dopamine interaction resulting in Fos production in the limbic system and basal ganglia of the mouse brain (75) 271
Reddy, S., see Scammell, J.G. (75) 8
Redman, C., see Rife, T.K. (75) 225
Renbäck, K., Inoue, M., Yoshida, A., Nyberg, F. and Ueda, H.
Vzg-1/lysophosphatidic acid-receptor involved in peripheral pain transmission (75) 350
Résibois, A., see Depaepe, V. (75) 76
Riera, L., see Goren, N. (75) 138
Rife, T.K., Xie, J., Redman, C. and Young, A.P.
The 5′2 promoter of the neuronal nitric oxide synthase dual promoter complex mediates inducibility by nerve growth factor (75) 225
Rodrigues, S.M., see Latimer, V.S. (75) 287
Romanovitch, A.E., see Plata-Salamán, C.R. (75) 248
Ross, R.A., see Scammell, J.G. (75) 8

Saito, S., see Kudo, M. (75) 172
Sands, S.A., Strong, R., Corbitt, J. and Morilak, D.A.
Effects of acute restraint stress on tyrosine hydroxylase mRNA expression in locus coeruleus of Wistar and Wistar-Kyoto rats (75) 1
Sarvey, J., see Valentine, G. (75) 337
Scammell, J.G., Reddy, S., Valentine, D.L., Coker, T.N., Nikolopoulos, S.N. and Ross, R.A.
Isolation and characterization of the human secretogranin II gene promoter (75) 8
Schaaf, M.J.M., Duurland, R., De Kloet, E.R. and Vreugdenhil, E.
Circadian variation in BDNF mRNA expression in the rat hippocampus (75) 342
Schaefer, A.L., see Weaver, S.A. (75) 46
Seo, K.H., see Lee, Z.H. (75) 237
Shibanuma, M., see Zhou, C.J. (75) 150
Shioda, S., see Zhou, C.J. (75) 150
Solórzano, G., see Talley, E.M. (75) 159
Sooy, K., see Gary, D.S. (75) 89
Spiess, J., see Radulovic, J. (75) 271
Steeves, T.D.L., see Zhu, H. (75) 303
Strong, R., see Sands, S.A. (75) 1
Sturani, E., see Ambrosini, A. (75) 54
Suzaki, H., see Kudo, M. (75) 172
Sweileh, W., Wenberg, K., Xu, J., Forsayeth, J., Hardy, S. and Loring, R.H.
Multistep expression and assembly of neuronal nicotinic receptors is both host-cell- and receptor-subtype-dependent (75) 293

Takahashi, J.S., see Zhu, H. (75) 303
Talley, E.M., Solórzano, G., Depaulis, A., Perez-Reyes, E. and Bayliss, D.A.
Low-voltage-activated calcium channel subunit expression in a genetic model of absence epilepsy in the rat (75) 159
Tapia-Arancibia, L., see Givalois, L. (75) 166
Terenius, L., see Broberger, C. (75) 25
Tessler, S., see Ingram, E.M. (75) 96
Thöny, B., see Depaepe, V. (75) 76
Tininini, S., see Ambrosini, A. (75) 54
Tokumaru, J., see Doi, T. (75) 105
Tsai, M.J., see Hung, H.-C. (75) 330
Turrin, N.P., see Plata-Salamán, C.R. (75) 248

Ueda, H., see Renbäck, K. (75) 350
Ueda, Y., see Doi, T. (75) 105
Urbanski, H.F., see Latimer, V.S. (75) 287

Valentine, D.L., see Scammell, J.G. (75) 8
Valentine, G., Chakravarty, S., Sarvey, J., Bramham, C. and Herkenham, M.
Fragile X (*fmr1*) mRNA expression is dif-

ferentially regulated in two adult models of activity-dependent gene expression (75) 337
Vana, A.M., see Naeve, G.S. (75) 185
Van Herp, F., see Holling, T.M. (75) 70
Ventura, F., see Goren, N. (75) 138
Verge, G., see Naeve, G.S. (75) 185
Vernet-der Garabedian, B., Lemaigre-Dubreuil, Y. and Mariani, J.
Central origin of IL-1β produced during peripheral inflammation: role of meninges (75) 259
Vreugdenhil, E., see Schaaf, M.J.M. (75) 342

Weaver, S.A., Schaefer, A.L. and Dixon, W.T.
The effects of mutated skeletal ryanodine receptors on calreticulin and calsequestrin expression in the brain and pituitary gland of boars (75) 46
Wenberg, K., see Sweileh, W. (75) 293
White, R.B., see Latimer, V.S. (75) 287
Whiteley, A., see Zhu, H. (75) 303
Willmore, L.J., see Doi, T. (75) 105
Willson, C.A., see Linnarsson, S. (75) 61
Wu, H.C., see Hung, H.-C. (75) 330

Xie, J., see Rife, T.K. (75) 225
Xu, J., see Sweileh, W. (75) 293

Yang, D.D., see Mielke, K. (75) 128
Yoshida, A., see Renbäck, K. (75) 350
Young, A.P., see Rife, T.K. (75) 225

Zhan-Poe, X., see Craft, C.M. (75) 198
Zhao, W., see D'Agata, V. (75) 345
Zhou, C.J., Kikuyama, S., Shibanuma, M., Hirabayashi, T., Nakajo, S., Arimura, A. and Shioda, S.
Cellular distribution of the splice variants of the receptor for pituitary adenylate cyclase-activating polypeptide (PAC_1-R) in the rat brain by in situ RT-PCR (75) 150
Zhu, H., LaRue, S., Whiteley, A., Steeves, T.D.L., Takahashi, J.S. and Green, C.B.
The *Xenopus Clock* gene is constitutively expressed in retinal photoreceptors (75) 303
Zippel, R., see Ambrosini, A. (75) 54
Zwiller, J., see Bernabeu, R. (75) 113